

# A Data Assimilation Testbed using KENDA and idealized COSMO Ensembles

Heiner Lange, George C. Craig

Hans-Ertel-Centre for Weather Research, Data Assimilation Branch

COSMO User Seminar  
Data Assimilation Session  
Offenbach, 17.03.2014

# Outline

## 1 idealized COSMO-KENDA


- Convective-scale Data Assimilation
- Nature Run and Synthetic Observations

## 2 Experiments: LETKF Cycling and Ensemble Forecasts

- Analysis Ensemble Convergence
- Forecast error growth
- Results, Conclusions, Outlook

# Thunderstorm Prediction


Large-scale phenomena (e.g. low-pressure systems) more predictable than small scales (convection, thunderstorms)



$\Delta t = 1 - 3 \text{ h} \rightarrow$  difficult predictability for  $\rightarrow \Delta x = 20 \text{ km}$

# Thunderstorm Prediction


Large-scale phenomena (e.g. low-pressure systems) more predictable than small scales (convection, thunderstorms)



$\Delta t = 1 - 3 \text{ h} \rightarrow$  difficult predictability for  $\rightarrow \Delta x = 20 \text{ km}$

# Thunderstorm Prediction


Large-scale phenomena (e.g. low-pressure systems) more predictable than small scales (convection, thunderstorms)



$\Delta t = 1 - 3 \text{ h} \rightarrow$  difficult predictability for  $\rightarrow \Delta x = 20 \text{ km}$

# Thunderstorm Prediction


Large-scale phenomena (e.g. low-pressure systems) more predictable than small scales (convection, thunderstorms)



$\Delta t = 1 - 3 \text{ h} \rightarrow$  difficult predictability for  $\rightarrow \Delta x = 20 \text{ km}$

# Thunderstorm Prediction


Large-scale phenomena (e.g. low-pressure systems) more predictable than small scales (convection, thunderstorms)



$\Delta t = 1 - 3 \text{ h} \rightarrow$  difficult predictability for  $\rightarrow \Delta x = 20 \text{ km}$

# Thunderstorm Prediction


Large-scale phenomena (e.g. low-pressure systems) more predictable than smalls scales (convection, thunderstorms)



$\Delta t = 1 - 3 \text{ h} \rightarrow$  difficult predictability for  $\rightarrow \Delta x = 20 \text{ km}$

# Thunderstorm Prediction

Large-scale phenomena (e.g. low-pressure systems) more predictable than small scales (convection, thunderstorms)



$\Delta t = 1 - 3 \text{ h} \rightarrow$  difficult predictability for  $\rightarrow \Delta x = 20 \text{ km}$

# Ensemble Data Assimilation

## Ensemble Forecasting and Data Assimilation:

- chaotic nature of atmosphere taken into account

# Ensemble Data Assimilation

## Ensemble Forecasting and Data Assimilation:

- chaotic nature of atmosphere taken into account
- ensemble forecasts more useful than deterministic forecasts:
  - ensemble mean represents state with highest probability
  - ensemble spread gives uncertainty information

# Ensemble Data Assimilation

## Ensemble Forecasting and Data Assimilation:

- chaotic nature of atmosphere taken into account
- ensemble forecasts more useful than deterministic forecasts:
  - ensemble mean represents state with highest probability
  - ensemble spread gives uncertainty information
- decision of DWD for operation of KENDA:
  - Kilometre-scale ENsemle Data Assimilation
  - Local Ensemble Transform Kalman Filter (LETKF) coupled with convection-permitting German forecast model  
COSMO-DE

# Data Assimilation of Convection

## Experiments for Ensemble Data Assimilation of Convection:

- **Perfect model** experiments (with nature run) to understand LETKF filter algorithms
- **Convection only**: no large scales
- **Simulated Dopper Radar** as realistic observation system

# Nature Run and Ensemble

COSMO model setup (based on Daniel Leuenberger's work)

Domain:  $198 \times 198 \times 50$  gridpoints  
periodic lateral boundaries conditions

Resolution: 2 km horizontally



Initial state: Horizontally homogenous sounding,  
 $\text{CAPE} = 2200 \frac{\text{J}}{\text{kg}}$ ,  
random white noise on T (0.02 K) and W ( $0.02 \frac{\text{m}}{\text{s}}$ )  
in the boundary layer

Model: Full COSMO physics with active radiation scheme



Forecast: 8 hour spinup until convection evolves:

- long-lived cells, lifetime  $\geq 6$  h
- horizontal position *fully random* in ensemble



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 



## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 


## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 


## 1 km Resolution vs. 2 km

 $\Delta x = 2 \text{ km}$  $\Delta x = 1 \text{ km}$ 


# Synthetic Observations

20 UTC: Synthetic Observations at  $z = 9776$  m


# Synthetic Observations

20 UTC: Synthetic Observations at  $z = 514$  m

## LETKF-Flowchart



# LETKF - How it works



# LETKF-Setup 1/2

## Assimilation setup

- 50 member ensemble (perfect model)
- Python-simulated observations of *radial wind* and (*no*)-reflectivity
- periodic LETKF-solution
- Python-cycling environment

# LETKF-Setup 1/2

## Assimilation setup


- 50 member ensemble (perfect model)
- Python-simulated observations of *radial wind* and (*no*)-reflectivity
- periodic LETKF-solution
- Python-cycling environment
- 3 hours cycled assimilation
- 3 hours ensemble forecast

# LETKF-Setup 2/2


## Assimilation settings

- Observation Resolution 2 and 8 km (superobservations SO)
- LETKF coarse grid on 2 and 8 km (coarse grid CG)
- Assimilation interval 5 and 20 minutes (20)
- Horizontal localization 8 and 32 km (L8 and L32)


# Assimilation Results: Nature vs. Analysis Ensemble Means




# Assimilation Results: Nature vs. Analysis Ensemble Means



# Assimilation Results: Nature vs. Analysis Ensemble Means




# Analysis Members L8




# Analysis Members L32SOCG20


L32SOCG20 Analysis,  $t = 17$  UTC



## Forecast error growth (displacement of storms)



# Forecast Results: Nature vs. Forecast Ensemble Means



# Results and Conclusions

## Assimilation Results

- Observation resolution of  $4 \times \Delta x_{model}$  sufficient for analysis
- Update of temperature through covariances: problematic!
- Less noise introduced with  $\Delta t_{ass} = 20$  minutes

# Results and Conclusions

## Assimilation Results

- Observation resolution of  $4 \times \Delta x_{model}$  sufficient for analysis
- Update of temperature through covariances: problematic!
- Less noise introduced with  $\Delta t_{ass} = 20$  minutes

## Forecast Results

- Very fast displacement error growth
- Bias issues with spurious convection
- Forecast quality limited by:
  - Predictability of convection
  - Dynamical imbalance of initial states (= LETKF analyses)

# Outlook 1/2

## Idealized KENDA

- Creative sandbox to study assimilation properties:
  - arbitrary types of convection (idealized COSMO)
  - free choice of ensemble spread / biases / model error
  - free choice of observation types, coverage and errors  
(very simple Python operator)
- KENDA-in-a-box: One Python cycling script  
(namelist-controlled) steers everything

# Outlook 2/2

## Further idealized studies

- Stability and imbalance in convective DA due to
  - strong ensemble convergence (observation errors and scale)
  - biases from strong localization
  - nonlinear observation operators
- Assimilation of convection in bad environmental soundings