

# Raytracing Berlin Neukölln




Berlin data from ([daten.berlin.de](http://daten.berlin.de))

## Milestones:

- Use environment data from project partners (e.g. Wieke Heldens, DLR)

# Raytracing Berlin Neukölln



Berlin data from ([daten.berlin.de](http://daten.berlin.de))

## Milestones:

- ▶ Use environment data from project partners (e.g. Wieke Heldens, DLR)
- ▶ Tessellation to get a raytraceable mesh (from CityGML data format to triangles)

# Raytracing Berlin Neukölln



Computations done with RayLi and libRadtran  
(Library for Radiative Transfer, [libradtran.org](http://libradtran.org))

## Milestones:

- ▶ Use environment data from project partners (e.g. Wieke Heldens, DLR)
- ▶ Tessellation to get a raytraceable mesh (from CityGML data format to triangles)
- ▶ Investigate 3D radiative effects in urban environments (including clouds, fog, and pollutants)

# Raytracing Berlin Neukölln




Computations done with RayLi and libRadtran  
(Library for Radiative Transfer, [libradtran.org](http://libradtran.org))

## Milestones:

- ▶ Use environment data from project partners (e.g. Wieke Heldens, DLR)
- ▶ Tessellation to get a raytraceable mesh (from CityGML data format to triangles)
- ▶ Investigate 3D radiative effects in urban environments (including clouds, fog, and pollutants)



# Raytracing Berlin Neukölln




Computations done with RayLi and libRadtran  
(Library for Radiative Transfer, [libradtran.org](http://libradtran.org))

## Milestones:

- ▶ Use environment data from project partners (e.g. Wieke Heldens, DLR)
- ▶ Tessellation to get a raytraceable mesh (from CityGML data format to triangles)
- ▶ Investigate 3D radiative effects in urban environments (including clouds, fog, and pollutants)

# Raytracing Berlin Neukölln




Computations done with RayLi and libRadtran  
(Library for Radiative Transfer, [libradtran.org](http://libradtran.org))

## Milestones:

- ▶ Use environment data from project partners (e.g. Wieke Heldens, DLR)
- ▶ Tessellation to get a raytraceable mesh (from CityGML data format to triangles)
- ▶ Investigate 3D radiative effects in urban environments (including clouds, fog, and pollutants)


# Raytracing Berlin Neukölln



Computations done with RayLi and libRadtran  
(Library for Radiative Transfer, [libradtran.org](http://libradtran.org))

## Milestones:

- ▶ Use environment data from project partners (e.g. Wieke Heldens, DLR)
- ▶ Tessellation to get a raytraceable mesh (from CityGML data format to triangles)
- ▶ Investigate 3D radiative effects in urban environments (including clouds, fog, and pollutants)



Computations done with RayLi and libRadtran  
(Library for Radiative Transfer, [libradtran.org](http://libradtran.org))

## Milestones:

- ▶ Use environment data from project partners (e.g. Wieke Heldens, DLR)
- ▶ Tessellation to get a raytraceable mesh (from CityGML data format to triangles)
- ▶ Investigate 3D radiative effects in urban environments (including clouds, fog, and pollutants)
- ▶ Introduce complex surfaces into the TenStream (LES radiation scheme)